Ve401 Probabilistic Methods in Engineering

Summer 2017 — Assignment 2

Date Due: 12:10 PM, Wednesday, the 31st of May 2017

This assignment has a total of (26 Marks).

Exercise 2.1 Discrete Uniform Distribution

A discrete random variable is said to be *uniformly distributed* if it assumes a finite number of values with each value occurring with the same probability. If we consider the generation of a single random digit taken from $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$, then Y, the number generated, is uniformly distributed with each possible digit occurring with probability 1/10. In general, the density for a uniformly distributed random variable is given by

$$f(x) = \frac{1}{n},$$
 $x = x_1, \dots, x_n,$ $n \in \mathbb{N}.$

- i) Find the moment-generating function for a discrete uniform random variable. (2 Marks)
- ii) Use the moment-generating function to find E[X] and Var[X]. (2 Marks)
- iii) Find the mean and variance for the random variable Y, the number obtained when a random digit generator is activated once.

(2 Marks)

Exercise 2.2 Uniqueness of Moment Generating Functions - Simple Case

Suppose that two discrete random variables (X, f_X) and (Y, f_Y) both take on values $\{0, \ldots, n\}$, $n \in \mathbb{N}$. Suppose that the moment-generating functions are equal in some neighborhood of zero, i.e., there exists some $\varepsilon > 0$ such that

 $m_X(t) = m_Y(t)$ for all $x \in (-\varepsilon, \varepsilon)$.

Show that $f_X(x) = f_Y(x)$ for x = 0, ..., n. (4 Marks)

Exercise 2.3

- i) Find the moment-generating function for a binomial random variable X with parameters n and p. (2 Marks)
- ii) Use the moment-generating function to find E[X] and Var X. (2 Marks)

Exercise 2.4 Drawing until First Success in the Hypergeometric Setting

i) Let (X, f_X) be a discrete random variable taking on only positive values, i.e., ran $X \subset \mathbb{N}$. Show that

$$\mathbf{E}[X] = \sum_{x=0}^{\infty} P[X > x].$$

(2 Marks)

ii) A box contains N balls, of which r are red and N - r are black. Balls are drawn from the box until a red ball is drawn. Show that the expected number of draws is

$$\frac{N+1}{r+1}.$$

Hint: $\sum_{x=0}^{r} \binom{N-r+x}{N-r} = \binom{N+1}{N-r+1}$ (4 Marks)



Exercise 2.5 Sums of Independent Discrete Random Variables

Two discrete random variables X and Y are said to be independent if

$$P[X = x \text{ and } Y = y] = P[X = x] \cdot P[Y = y] \qquad \text{for any } x \in \operatorname{ran} X \text{ and } y \in \operatorname{ran} Y.$$

i) Let Z = X + Y, where X and Y are assumed to be independent. Show that

$$P[Z=z] = \sum_{x+y=z} P[X=x] \cdot P[Y=y]$$

using the formula for total probability. (3 Marks)

ii) Show that the sum of two independent geometric random variables follows a Pascal distribution with r = 2.

(3 Marks)